

Du Pont (UK) Limited, Crop Protection Products - 4th Floor,
Kings Court, London Road, STEVENAGE, Herts. SG1 2NG -
Tel: 0800 689 8899 (UK): +44 (0) 689 8899 or
email: ukhotline@corvea.com

24 hour Emergency Tel: +44 (0) 161 884 1235 (UK) /
+353 76 680 5288 (IRL)

National Poisons information tel:

111 (UK), or 01 809 2166 (Ireland)

PROTECT FROM FROST.

SHAKE WELL BEFORE USE FOR PROFESSIONAL USE ONLY

To access the Safety Data Sheet for this product scan the QR
code or use the weblink below:

<https://www.corvea.co.uk/content/dam/dpagco/corvea/eu/gb/en/files/sds/TALIUS-SDS.pdf>

<https://www.corvea.ie/content/dam/dpagco/corvea/eu/ie/en/files/sds/TALIUS-IRL-SDS.pdf>



The Voluntary Initiative
This label is compliant with
the CPA Voluntary Initiative
Guidance (UK only)



A60007856

Manufactured in E.U.

Batch N°.: See top of pack



Talius™

FUNGICIDE

GROUP

13

FUNGICIDE

MAPP No 12752
PCS No 04604



An emulsifiable concentrate formulation containing 200 g/litre proquinazid for control of powdery mildew in wheat, barley, oats, winter rye, spring rye (Ireland only) and triticale.

RISK AND SAFETY INFORMATION

TALIUS™

Fungicide – contains 200 g/litre proquinazid

Causes skin irritation.

Causes serious eye damage.

Suspected of causing cancer.

Very toxic to aquatic life with long lasting effects.

Obtain special instructions before use.

Wear protective gloves/protective clothing / eye protection/face protection.

IF ON SKIN: Wash with plenty of soap and water.

IF IN EYES: Rinse cautiously with water for several minutes.

Remove contact lenses, if present and easy to do. Continue rinsing.

IF exposed or concerned: Get medical advice/attention.

Immediately call a POISON Centre or doctor/physician.

Take off contaminated clothing and wash before reuse.

Collect spillage.


Store locked up.

Dispose of contents / container to a licensed hazardous-waste disposal contractor or collection site except for triple rinsed empty clean containers which can be disposed of as non-hazardous waste.

To avoid risks to human health and the environment, comply with the instructions for use.

MAPP No 12752 / PCS No 04604

DANGER



Nett Contents: 1 litre e

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work (UK only).

SAFETY PRECAUTIONS

Operator protection

RISK OF SERIOUS DAMAGE TO EYES

IRRITATING TO SKIN

IN CASE OF CONTACT WITH EYES, RINSE IMMEDIATELY with plenty of water and seek medical advice.

IF YOU FEEL UNWELL, seek medical advice immediately (show the label where possible)

WHEN USING DO NOT EAT, DRINK OR SMOKE

(UK only) Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS), SUITABLE PROTECTIVE GLOVES,

FACE PROTECTION (FACESHIELD) when handling the concentrate.

WEAR SUITABLE PROTECTIVE GLOVES when handling contaminated surfaces.

(UK only) However, engineering controls may replace personal protective equipment if a COSHH assessment shows that they provide equal or higher standard of protection.

Environmental protection

Do not contaminate water with the product or its container (Do not clean application equipment near surface water / Avoid contamination via drains from farmyards and roads).

(UK only) To protect aquatic organisms respect an unsprayed buffer zone to surface water bodies in line with LERAP requirements. **DO NOT ALLOW DIRECT SPRAY** from horizontal boom sprayers to fall within 5 m of the top of the bank of a static or flowing waterbody, unless a Local Environmental Risk Assessment for Pesticides (LERAP) permits a narrower buffer zone, or within 1 m of the top of a ditch which is dry at the time of application. **DO NOT ALLOW DIRECT SPRAY** from hand-held sprayers to fall within 1 m of the top of a bank of a static or flowing waterbody. Aim spray away from water.

This product qualifies for inclusion within the Local Environmental Risk Assessment for Pesticides (LERAP) scheme. Before each spraying operation from a horizontal boom sprayer, either a LERAP must be carried out in accordance with CRD's published guidance or the statutory buffer zone must be maintained. The results of the LERAP must be recorded and kept available for three years.



(Ireland only) To protect aquatic organisms respect an unsprayed buffer zone of 5m to surface water bodies

Storage and disposal

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDING STUFFS

KEEP OUT OF REACH OF CHILDREN

KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place

WASH OUT CONTAINER THOROUGHLY, empty washings into spray tank and dispose of safely

EMPTY CONTAINER COMPLETELY and dispose of safely

DO NOT RE-USE CONTAINER for any purpose

™ © Trademarks of Corteva Agriscience
and its affiliated companies.



3 576960 031165



DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

Restrictions

TALIUS™ should not be applied to any crop suffering from stress as a result of drought, waterlogging, low temperatures, nutrient or lime deficiency or other factors reducing crop growth.

- Avoid application in either frosty or hot, sunny weather conditions.
- Up to 2 treatments may be applied per crop, however the second application should be tankmixed with an alternate mode of action mildewicide (see Timing).
- There is a possibility that some crops, particularly oilseed rape and other brassicas, may suffer adverse effects from TALIUS™, care should therefore be taken to avoid drift onto non-target crops (UK only).

Disease Control

TALIUS™ controls Powdery mildew in cereals. Application should be made preventatively before the disease has established in the crop. If mildew has actively spread to new growth TALIUS™ should be tankmixed with a curative, alternate mode of action mildewicide.

Crops

TALIUS™ can be used on all varieties of wheat, barley, oats, winter rye, spring rye (Ireland only) and triticale.

Timings

TALIUS™ can be applied at any time before full flowering (GS 65) on wheat or before beginning of heading (GS 49) on barley, oats, winter rye, spring rye (Ireland only) and triticale. Up to a maximum of 2 treatments may be made. The first treatment should be applied preventatively at the onset of disease development before its spread to new growth. If a second treatment is required, this may be applied after 4 – 6 weeks and must be tankmixed with an alternate mode of action mildewicide (see Compatibility).

Volume and application

Apply TALIUS™ in 150 - 400 litres of water per hectare as a medium quality spray (as defined by BCPC). A spray pressure of 2 – 3 Bar is recommended. When applying TALIUS™, care should be taken not to overlap spray swaths.

Dose

Apply TALIUS™ at 0.25 litres/hectare.

Following crops

There are no restrictions on crops that may be grown following harvest of a cereal crop treated with Talius.

Resistance

TALIUS™ contains a quinazolinone group fungicide and must be considered to be in the same group for resistance management purposes as quinolines. There is a significant risk of widespread resistance occurring in Powdery mildew populations and failure to follow resistance management action may result in reduced levels of disease control. Development of resistance can be avoided or delayed by alternating or mixing products having an alternate mode of action (see Timing). Use TALIUS™ as part of an Integrated Crop Management Strategy (ICM) incorporating other methods of control, including where appropriate other fungicides with a different mode of action.

When fungicides with the same mode of action are used repeatedly over several years in the same field, naturally occurring less sensitive strains may survive, propagate and become dominant in that field. A pathogen is considered resistant to a fungicide if it survives a correctly applied treatment at a recommended dose and timing, under normal weather conditions.

Mixing

Half-fill the tank with clean water, add the required amount of TALIUS™ and continue agitation while topping up the tank and while spraying. Wash out container and add washings to the spray tank.

Compatibility

For resistance management purposes, TALIUS™ may also be tankmixed with alternate mode of action mildewicides. Products should only be tank-mixed if each product can be applied within the label

